S.C.E.N.T.

Southern California Ear, Nose and Throat 433 E. Wardlow Rd. Long Beach, CA 90807

Phone: 562.427.0550 Fax: 562.988.8899

Post-Operative Instructions for Frenuloplasty-Release of Tongue Tie

Procedure:

A piece of the mucosa (skin in the mouth), and sometimes a tight muscle under the tongue, has been realigned to give your child's tongue better movement. A small suture (stitch) was used to repair the area. It will dissolve on its own and will not need to be removed.

INSTRUCTIONS FOR HOME:

Expectations: Most children have a very little discomfort. Bleeding or swelling is not expected so please call us if either of these problems occurs.

Diet: A normal diet may be resumed after discharge

Activity: Normal activities may be resumed after discharge

Medications: Please give Tylenol if your child experiences discomfort

DO NOT USE ASPIRIN OR IBUPROFIN PRODUCTS SUCH AS ASPERGUN, ALEVE, MOTRIN, OR ADVIL AS THEY CAN CONTRIBUTE TO BLEEDING. AVOID SUCH HERBALS AS GINSENG, GINGKO, OR GARLIC SUPPLEMENTS

Follow-Up: Schedule the patient for a follow-up appointment in our office 3-6 weeks after surgery.

If you have any questions, problems, or need to schedule (or reschedule) an appointment, please contact our office at 562.427.0550